



PrudentMD Aesthetics

5 Dundee Avenue, Iselin, NJ 08830

Ph: (732) 404-0044 | Fax: (732) 218-3933 | www.prudentmd.com

Botox / Xeomin - information

Post-procedure:

- After injection of very small amount of Botox cosmetic, it results in relaxation of muscle and fine lines. This response usually appears 2-6 days post procedure.
- Stay erect and do not manipulate the area for at least 4 hours post-procedure.
- It is recommended that you visit the physician again after 1 week of injection for post-treatment evaluation.
- You may experience headache, bruising, pain at injection site.
- Rarely asymmetry, twitching, numbness, temporary drooping of eyelids or eyebrows may be seen.

General guidelines:

- Pregnant, breast-feeding females must not get the procedure done.
- Contact the physician at 732-404-0044 for any concern that may develop post-procedure.
- Allergic reaction can rarely occur just like any other product.
- Individuals with peripheral motor neuropathic disease, amyotrophic lateral sclerosis, myasthenia gravis, Lambert-Eaton syndrome must not get Botox cosmetic procedure done.
- Botox cosmetic is not recommended in children younger than 18 years of age.
- Botox cosmetic must not be used for individuals with allergy to Botox, Dysport, Myobloc and for individuals with recent skin injection at planned injection site.
- Please inform doctor if you have taken Botox, dysport or myobloc, fillers in last 6 months; cold medicine, sleeping pill, muscle relaxant, allergy medicine etc. in past 1 week.
- Botox cosmetic is best at treating dynamic facial lines caused by facial muscle activity. Lines present at rest may or may not improve.
- Typically, wrinkles will reappear in 3-5 months. At this point, repeat treatment will relax the lines again.
- Rarely injection produces unsatisfactory result. Therefore, it is recommended to have 1 week follow up post-procedure.